

National Guidelines for Palliative Care

**First edition released 28 June 2014,
Revised on 20th Jan 2015**

Contents

Foreword	3
Introduction.....	4
WHO Definition of Palliative Care.....	5
Definition of Other Terms.....	6
How to Use the Guidelines	8
Overview of the Guidelines	9
Domain 1: Patient Care.....	11
<i>Guideline 1</i> – Timely Identification.....	11
<i>Guideline 2</i> – Reducing Barriers to Care.....	13
<i>Guideline 3</i> – Coordinated Care	15
<i>Guideline 4</i> – Holistic Assessment and On-going Care Planning	17
<i>Guideline 5</i> – Advance Care Planning.....	19
<i>Guideline 6</i> – Patient-Centred Care.....	20
<i>Guideline 7</i> – Care in the Last Days of Life	22
Domain 2: Family and Caregiver Support	24
<i>Guideline 8</i> – Caregiver Support.....	24
<i>Guideline 9</i> – Bereavement Care.....	26
Domain 3: Staff and Volunteer Management.....	Error! Bookmark not defined.
<i>Guideline 10</i> – Qualified Staff and Volunteers	28
<i>Guideline 11</i> – Staff and Volunteer Self-Care	30
Domain 4: Safe Care	31
<i>Guideline 12</i> – Access to and Use of Opioids	31
<i>Guideline 13</i> – Clinical Quality Improvement	33

Foreword

In 2010, the Ministry of Health of Singapore commissioned the Lien Centre for Palliative Care to formulate a National Strategy for Palliative Care in consultation with key stakeholders in the healthcare system. In the report released in 2011, one of the goals of the national strategy called for the development of local standards in palliative care. The Standards Development Subgroup of the National Strategy for Palliative Care Implementation Taskforce is now pleased to provide the local community with the inaugural edition of the National Guidelines for Palliative Care.

The integration of palliative care within the healthcare system means that one of the major challenges faced by the workgroup is the need for the guidelines to be applicable across different care settings. Hence, the workgroup has divided service providers into different classes according to the nature and scope of their work in providing palliative care. The indicators derived are based on the level of care expected of each class of service provider.

A participatory process was used in the development of these guidelines. Draft guidelines were developed by the workgroup after an initial literature review and consultation with key stakeholders on service gaps and challenges. With each revision, the views of focal persons were canvassed. Finally, tools were suggested which can be adapted for each provider's use in the use of the guidelines.

The guidelines recognise that individual patients have different needs at different phases of their illness and service providers should be responsive to patients' changing needs. In addition, families and carers need support during the patient's life and in bereavement. It also recognises the importance of training and self-care for staff and the important role of volunteers in palliative care.

It is hoped that these guidelines will provide guidance in the delivery of high-quality care for the terminally ill, minimise gaps in service, improve the quality of training as well as ensure support for all staff and volunteers serving in this field.

This milestone would not have been reached without the contribution of many. We extend our grateful thanks to members of the workgroup who helped to put this document together, the secretariat for their support and all the healthcare professionals who gave us the benefit of their experience and thoughtful comments.

A/Prof Pang Weng Sun

Chairman

National Strategy for Palliative Care Implementation Taskforce

Dr Angel Lee

Chairman

Standards Development Subgroup

Vice-Chairman

National Strategy for Palliative Care Implementation Taskforce

Introduction

Palliative care in Singapore has come a long way since its humble beginnings in the 1980s. With the ageing population and increasing incidence of cancer and chronic diseases, the demand for palliative care will continue to rise in the future. The National Strategy for Palliative Care, accepted by the Ministry of Health in 2012, outlined ten key goals for palliative care in Singapore. Among them was that there should be local standards of care to ensure the delivery of good quality palliative care.

Patients with life-limiting illnesses and their families have numerous physical, psychosocial and spiritual needs as they approach the end of life. Palliative care aims to meet these needs in a holistic manner. The World Health Organization (WHO) defines palliative care as “an approach that improves the quality of life of patients and their families facing the problem associated with life-threatening illness, through the prevention and relief of suffering by means of early identification and impeccable assessment and treatment of pain and other problems, physical, psychosocial and spiritual”¹.

These patients and their families receive care from a multitude of healthcare providers across the healthcare spectrum, in settings ranging from primary care in the community to palliative care services in hospices and hospitals. The holistic approach described by WHO should be practised by **all providers** who care for such patients and their families, regardless of setting.

As outlined in the National Strategy for Palliative Care, this first edition of the Guidelines for Palliative Care in Singapore aims to articulate a vision for high quality palliative care, through providing evidence-based guidelines for the holistic approach described above.

It is recognised that the different groups of providers across the healthcare spectrum have differing roles in the provision of palliative care. As such, three groups of providers have been identified, and specific indicators have been described for each of these groups. This will be elaborated on in subsequent sections (see *Definition of Other Terms*).

It is envisioned that in this manner, the Guidelines will promote a whole-of-sector approach to providing accessible, high-quality palliative care in Singapore, so that all who suffer from life-limiting illnesses may live their last days in peace, comfort and dignity.

¹ WHO definition of palliative care (accessed 15 Mar 2013), available at <http://www.who.int/cancer/palliative/definition/en/>

WHO Definition of Palliative Care

The World Health Organisation describes palliative care as:

“...an approach that improves quality of life of patients and their families facing the problem associated with life-threatening illness, through the prevention and relief of suffering by means of early identification and impeccable assessment and treatment of pain and other problems, physical, psychological and spiritual.”

Palliative care:

- provides relief from pain and other distressing symptoms;
- affirms life and regards dying as a normal process;
- intends neither to hasten or postpone death;
- integrates psychological and spiritual aspects of patient care;
- offers a support system to help patients live as actively as possible until death;
- offers a support system to help the family cope during the patient’s illness and in bereavement;
- uses a team approach to address the needs of patients and their families, including bereavement counselling if indicated;
- will enhance quality of life, and may also positively influence the course of illness;
- is applicable early in the course of illness, in conjunction with other therapies that are intended to prolong life, such as chemotherapy or radiation therapy, and includes those investigations needed to better understand and manage distressing clinical complications.²

² WHO definition of palliative care (accessed 15 Mar 2013), available at <http://www.who.int/cancer/palliative/definition/en/>

Definition of Other Terms

Life-limiting illness refers to illnesses where there is little or no hope of cure, and it is expected that death will be a direct consequence of the illness. This includes illnesses of both malignant and non-malignant nature.

Patient refers to the primary recipient of palliative care.

Caregiver refers to a person, often but not necessarily a family member, who undertakes to provide for the needs of the patient and who may take on additional technical tasks in the process, e.g. administration of medicines. The *primary caregiver* is the primary support person for the patient. This does not include members of the healthcare team.

Complex needs may derive from the patient, carer or health care team and the help required may be intermittent or continuous, depending on the level of need and rate of disease progression. Examples of complex levels of need include³:

- a) Physical symptoms - uncontrolled or complicated symptoms, specialised nursing requirements, complex mobility or functioning issues.
- b) Psychological - uncontrolled anxiety or depression, cognitive or behavioural issues.
- c) Social - complex situations involving children, family or carers, finance issues, communication difficulties and patients with special needs.
- d) Spiritual - unresolved issues around self-worth, loss of meaning and hope, requests for euthanasia, unresolved religious or cultural issues.
- e) Ethical - conflicting interests involving ethical principles that impinge on decision-making by patient, family or care team.

The Guidelines identify three groups of healthcare providers: **Class A**, **Class B** and **Class C** providers.

- **Class A providers** are those whose substantive work is not in caring for patients with life-limiting illnesses, but who will encounter them in the course of work. These include general practitioners in the community, and doctors, nurses and allied health staff in restructured and community hospitals.
- **Class B providers** are those who routinely care for a substantive number of patients with life-limiting illness. These include staff of chronic disease management programmes, intensive care units, specialist cancer units, geriatric units, home care providers and nursing homes.
- **Class C providers** are those who care solely for patients with life-limiting illness. These include palliative care teams in private, restructured and community hospitals, inpatient hospices and hospice home care and hospice day care providers.

³ North Yorkshire and York Palliative Care Group. Eligibility criteria for Specialist Palliative Care Services. 2005.

All classes of providers should utilise the palliative care approach in managing patients with life-limiting illness who are under their care. Notwithstanding this, it is recognised that the three groups have different roles in the provision of palliative care. For example, Class C providers manage patients whose needs exceed the capabilities of Class A and Class B providers, and also provide consultative support to other providers. Class A and Class B providers manage patients within their capabilities, but are responsible for referring patients and their families to Class C providers where appropriate. Therefore, different quality indicators have been outlined for each group, reflecting these differences in roles.

How to Use the Guidelines

The Guidelines are presented in three parts:

Guideline – the specific guiding statement

Rationale – the rationale behind adopting the Guideline

Quality Indicators – the specific, measurable indicators that reflect the Guideline in practice. The indicators are organised by the three groups of healthcare providers outlined in the preceding section.

Tools – examples of instruments, methods and other resources helpful in implementing or meeting the Guideline.

Overview of the Guidelines

Domain 1: Patient Care

Guideline 1 – Timely Identification

People approaching the end of life are identified in a timely manner.

Guideline 2 – Reducing Barriers to Care

Palliative care is available for all people based on clinical need, regardless of diagnosis, age, gender, financial ability, ethnic and cultural background, and care setting.

Guideline 3 – Coordinated Care

Care is delivered in a coordinated manner that ensures continuity of care across settings and over time.

Guideline 4 – Holistic Assessment and On-going Care Planning

Holistic assessment and on-going care planning are implemented in an interdisciplinary manner to meet the changing needs and wishes of patients, caregivers and families.

Guideline 5 – Advance Care Planning

All patients approaching the end of life have access to Advance Care Planning (ACP).

Guideline 6 – Patient-Centred Care

Patients have unique needs and preferences, which may differ depending on their cultural background. The patient's quality of life is improved by care that is customised to their unique physical, cultural, spiritual etc. needs.

Guideline 7 – Care in the Last Days of Life

Care is taken to fulfil the needs of patients in the last days of life, as well as that of their caregivers and families.

Domain 2: Family and Caregiver Support

Guideline 8 – Caregiver Support

Caregivers of patients with life-limiting illness face significant stress in their roles, and their own practical and emotional needs need to be supported.

Guideline 9 – Bereavement Care

Family members affected by a death are offered timely bereavement support appropriate to their needs and preferences.

Domain 3: Staff and Volunteer Management

Guideline 10 – Qualified Staff and Volunteers

Care for those approaching the end of life is delivered by staff and volunteers (where applicable) with the appropriate qualifications and skill mix for the level of service offered, and who demonstrate ongoing participation in training and development.

Guideline 11 – Staff and Volunteer Self-Care

Staff and volunteers reflect on practice, maintain effective self-care strategies and have access to support in dealing with the psychological stress associated with working among the terminally ill and bereaved.

Domain 4: Safe Care

Guideline 12 – Access to and Use of Opioids

Patients at the end of life should have access to opioids for symptom control, with guidelines and processes in place to ensure safe and effective use.

Guideline 13 – Clinical Quality Improvement

The service is committed to improvement in clinical and management practices.

Domain 1: Patient Care

Guideline 1 – Timely Identification

People approaching the end of life are identified in a timely manner.

Rationale Failure to identify patients with life-limiting illnesses may lead to inappropriate care. Timely identification with appropriate needs assessment helps service providers to meet patients’ needs and preferences in a more meaningful way.¹⁻¹²

Indicators

A + B

1.1 There is evidence of a system in place to identify people approaching the end of life (i.e. likely to die within the next 12 months).

Tools

1	<p>General prognostication and needs identification tools:</p> <ul style="list-style-type: none"> a) Gold Standards Framework (GSF) Prognostic Indicator Guidance¹³ b) Centre to Advance Palliative Care (CAPC) consensus criteria¹⁴ c) Supportive and Palliative Care Indicators Tool (SPICT™)¹⁵ 	-
2	<p>Disease-specific indicators of prognosis:</p> <ul style="list-style-type: none"> a) Heart failure: <ul style="list-style-type: none"> i. End of Life/ Palliative Education Resource Centre’s (EPERC) Fast Facts #143¹⁶ ii. Seattle Heart Failure Model¹⁷ b) COPD: <ul style="list-style-type: none"> i. EPERC Fast Facts #141¹⁸ ii. BODE scale¹⁹ c) Renal failure <ul style="list-style-type: none"> i. EPERC Fast Facts #191²⁰ d) Liver failure <ul style="list-style-type: none"> ii. EPERC Fast Facts #189²¹ e) Dementia: <ul style="list-style-type: none"> i. EPERC Fast Facts #150²² 	-

References:

1. McNamara BA, Rosenwax LK, Murray K, Currow DC. Early admission to community-based palliative care reduces use of emergency departments in the ninety days before death. *J Palliat Med* 2013;16(7):774-9.
2. Daveson BA, Alonso JP, Calanzani N, Ramsenthaler C, Gysels M, Antunes B, Moens K, Groeneveld EI, Albers G, Finetti S, Pettenati F, Bausewein C, Higginson IJ, Harding R, Deliens L, Toscani F, Ferreira PL, Ceulemans L, Gomes B; on behalf of PRISMA. Learning from the public: citizens describe the need to improve end-of-life care access, provision and recognition across Europe. *Eur J Public Health*. 2013 Jul 1. [Epub ahead of print]

3. Gaertner J, Weingärtner V, Wolf J, Voltz R. Early palliative care for patients with advanced cancer: how to make it work? *Curr Opin Oncol* 2013;25(4):342-52.
4. Glare PA. Early implementation of palliative care can improve patient outcomes. *J Natl Compr Canc Netw* 2013;11 Suppl 1:S3-9.
5. Bandieri E, Sichetti D, Romero M, Fanizza C, Belfiglio M, Buonaccorso L, Artioli F, Campione F, Tognoni G, Luppi M. Impact of early access to a palliative/supportive care intervention on pain management in patients with cancer. *Ann Oncol* 2012;23(8):2016-20.
6. Greer JA, Pirl WF, Jackson VA, Muzikansky A, Lennes IT, Heist RS, Gallagher ER, Temel JS. Effect of early palliative care on chemotherapy use and end-of-life care in patients with metastatic non-small-cell lung cancer. *J Clin Oncol* 2012;30(4):394-400.
7. Meier DE. Increased access to palliative care and hospice services: opportunities to improve value in health care. *Milbank Q.* 2011;89(3):343-80. doi: 10.1111/j.1468-0009.2011.00632.x.
8. Jones BW. The need for increased access to pediatric hospice and palliative care. *Dimens Crit Care Nurs* 2011;30(5):231-5.
9. Vassal P, Le Coz P, Herve C, Matillon Y, Chapuis F. Is the principle of equal access for all applied in practice to palliative care for the elderly? *J Palliat Med.* 2009;12(12):1089.
10. Morita T, Miyashita M, Tsuneto S, Sato K, Shima Y. Late referrals to palliative care units in Japan: nationwide follow-up survey and effects of palliative care team involvement after the Cancer Control Act. *J Pain Symptom Manage* 2009;38(2):191-6.
11. Temel JS, Greer JA, Muzikansky A, Gallagher ER, Admane S, Jackson VA, Dahlin CM, Blinderman CD, Jacobsen J, Pirl WF, Billings JA, Lynch TJ. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363(8):733-42.
12. Shaw KL, Clifford C, Thomas K, Meehan H. Review: improving end-of-life care: a critical review of the gold standards framework in primary care. *Palliat Med.* 2010 Apr;24(3):317-29.
13. National Gold Standards Framework Prognostic Indicator Guidance. Available online <http://www.goldstandardsframework.org.uk/content/uploads/files/General%20Files/Prognostic%20Indicator%20Guidance%20October%202011.pdf>. Last accessed 30th December 2013.
14. Weissman DE, Meier DE. Identifying patients in need of a palliative care assessment in the hospital setting: a consensus report from the center to advance palliative care. *J Palliat Med* 2011;14:17-23.
15. The University of Edinburgh, and NHS Lothian. SPICt. Available online <http://www.spict.org.uk/>. Last accessed 23rd January 2014.
16. Medical College of Wisconsin. #143 Prognostication in Heart Failure. Available online http://www.eperc.mcw.edu/EPERC/FastFactsIndex/ff_143.htm. Last accessed 18 February 2014.
17. University of Washington. Seattle Heart Failure Model. Available online <http://depts.washington.edu/shfm/>. Last accessed 23rd January 2014.
18. Medical College of Wisconsin. #141 Prognosis in End-Stage COPD. Available online http://www.eperc.mcw.edu/EPERC/FastFactsIndex/ff_141.htm. Last accessed 18 February 2014.
19. Celli BR, Cote CG, Marin JM, et al. The body-mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *N Eng J Med.* 2004; 350(10):1005-12.
20. Medical College of Wisconsin. #191 Prognostication in Patients Receiving Dialysis. Available online http://www.eperc.mcw.edu/EPERC/FastFactsIndex/ff_191.htm. Last accessed 18 February 2014.
21. Medical College of Wisconsin. #189 Prognosis in Decompensated Chronic Liver Failure. Available online http://www.eperc.mcw.edu/EPERC/FastFactsIndex/ff_189.htm. Last accessed 18 February 2014.
22. Medical College of Wisconsin. #150 Prognostication in Dementia. Available online http://www.eperc.mcw.edu/EPERC/FastFactsIndex/ff_150.htm. Last accessed 18 February 2014.

Domain 1: Patient Care

Guideline 2 – Reducing Barriers to Care

Palliative care is available for all people based on clinical need, regardless of diagnosis, age, gender, financial means, ethnic and cultural background, and care setting.

Rationale

All patients with life-limiting illnesses should be cared for by healthcare professionals using a palliative care approach. Patients with needs that exceed the resourced capabilities of the service provider should have access to palliative care services¹⁻¹¹. Similarly, where patients require expertise or care outside the scope of the palliative care service, the ability to call upon other services or providers will enhance the care of the patient.

Indicators

A + B	2.1	Patients are referred to palliative care services or providers in other fields (e.g. counselling support), should their needs exceed the resourced capabilities of the service provider.
	2.2	There is evidence of arrangements to ensure that people approaching the end of life, as well as their families and caregivers, know who to contact for advice.
C	2.3	The service provider triages and assigns priorities to all initial consult requests and ensures that care is delivered in a timely manner.
	2.4	The patient and family have access to specialist support 24 hours a day, seven days a week.
	2.5	The service provider has protocols for responding to palliative care emergencies or urgent needs.
	2.6	The service provider has formal links with specialists or providers in other fields to ensure access to expert advice and management of patients with specific needs in these areas.

Tools

1	Guidelines for referrals to palliative care/hospice services, or other specialists ¹²	-
2	Information for patients and families on scope of services provided by palliative care or hospice services and referral procedures ¹³	-
3	Protocols for responding to palliative care emergencies or urgent needs	-

References:

1. McVey P, McKenzie H, White K. A community-of-care: the integration of a palliative approach within residential aged care facilities in Australia. *Health Soc Care Community*. 2013 Nov 6. doi: 10.1111/hsc.12077. [Epub ahead of print]
2. Quill TE, Abernethy AP. Generalist plus specialist palliative care--creating a more sustainable model. *N Engl J Med* 2013;368(13):1173-5.
3. Reville B, Reifsnnyder J, McGuire DB, Kaiser K, Santana AJ. Education and referral criteria: impact on oncology referrals to palliative care. *J Palliat Med* 2013;16(7):786-9.
4. Addicott R. Delivering better end-of-life care in England: barriers to access for patients with a non-cancer diagnosis. *Health Econ Policy Law* 2012;7(4):441-54.

5. Nyatanga B. The pursuit of cultural competence: service accessibility and acceptability. *Int J Palliat Nurs* 2011;17(5):212-215.
6. Le BH, Watt JN. Care of the dying in Australia's busiest hospital: benefits of palliative care consultation and methods to enhance access. *J Palliat Med* 2010;13(7):855-60.
7. Lyckholm JJ, Coyne PJ, Kreutzer KO, Ramakrishnan V, Smith TJ. Barriers to effective palliative care for low-income patients in the late stages of cancer: A report of a study and strategies for defining and conquering the barriers. *Nurs Clin N Am* 2010;45(3):399-409.
8. Vassal P, Le Coz P, Herve C, Matillon Y, Chapuis F. Is the principle of equal access for all applied in practice to palliative care for the elderly? *J Palliat Med* 2009;12(12):1089.
9. Brumley R, Enguidanos S, Jamison P, Seitz R, Morgenstern N, Saito S, McIlwane J, Hillary K, Gonzalez J. Increased satisfaction with care and lower costs: results of a randomized trial of in-home palliative care. *J Am Geriatr Soc* 2007; 55 (7): 993-1000.
10. Birks T, Krikos D, McGowan C, Stone P. Is there a need for weekend face-to-face inpatient assessments by hospital specialist palliative care services? Evaluation of an out-of-hours service. *Palliat Med* 2011;25(3):278-283.
11. Kendall C, Jeffrey D. Out-of-hours specialist palliative care provision in an oncology centre: is it worthwhile? *Palliat Med* 2003;17(5):461-464.
12. Singapore Hospice Council. Who needs Hospice and Palliative Care? Available online. http://www.singaporehospice.org.sg/7.3_whoneedsit.htm. Last accessed 30th December 2013.
13. Singapore Hospice Council. Providers and Services in Singapore. Available online. http://www.singaporehospice.org.sg/7.4_providersandservices.htm. Last accessed 30th December 2013.

Domain 1: Patient Care

Guideline 3 – Coordinated Care

Care is delivered in a coordinated manner that ensures continuity of care across settings and over time.

Rationale

Patients may require care from various service providers at different stages of the illness. Poor coordination includes ineffective communication between services, inflexible responses to changes in need over time and fragmented care across different disciplines. The lack of coordination of care and services may increase the stress experienced by patients and their families, and lead to unnecessary resource utilisation (e.g. laboratory investigations, emergency department visits, admissions). Better coordination of care increases quality of life.¹⁻⁷

Indicators

All	3.1	There should be a primary provider coordinating the patient's care.
	3.2	Networks are established between different service providers, to facilitate the provision of seamless and holistic care for patients.
	3.3	The patient, caregivers and family are provided with clear written instructions on how to seek help if needed at any time, including after office hours.
	3.4	During transfers between different care settings, necessary patient information is provided to the receiving service provider.
	3.5	Where the patient's needs fall beyond the usual scope of service, for example personal care needs, referrals are made to other appropriate service providers to meet these needs.
	3.6	A plan is in place for the certification of death during and after office hours.
C	3.7	There is evidence of audits to ensure coordination of care.

Tools

1	Forms to aid staff in transfer of information: a. Hand-over forms when organisational boundaries are crossed (e.g. at clinic visits, referral to emergency departments)	-
2	Data monitoring and collection: a. Monitoring and audit of emergency department visits	-
3	Access to National Electronic Health Record (NEHR)	-

References:

1. Mason B, Epiphaniou E, Nanton V, Donaldson A, Shipman C, Daveson BA, Harding R, Higginson I, Munday D, Barclay S, Boyd K, Dale J, Kendall M, Worth A, Murray SA. Coordination of care for individuals with advanced progressive conditions: a multi-site ethnographic and serial interview study. *Br J Gen Pract* 2013;63(613):e580-8.
2. Husain A, Barbera L, Howell D, Moineddin R, Bezjak A, Sussman J. Advanced lung cancer patients' experience with continuity of care and supportive care needs. *Support Care Cancer* 2013;21(5):1351-8.
3. Morita T, Miyashita M, Yamagishi A, Akiyama M, Akizuki N, Hirai K, Imura C, Kato M, Kizawa Y, Shirahige Y, Yamaguchi T, Eguchi K. Effects of a programme of interventions on regional

comprehensive palliative care for patients with cancer: a mixed-methods study. *Lancet Oncol* 2013;14(7):638-46.

4. Engelhardt JB, McClive-Reed KP, Toseland RW, Smith TL, Larson DG, Tobin DR. Effects of a program for coordinated care of advanced illness on patients, surrogates, and healthcare costs: a randomized trial. *Am J Manag Care* 2006;12(2):93-100.
5. Aiken LS, Butner J, Lockhart CA, Volk-Craft BE, Hamilton G, Williams FG. Outcome evaluation of a randomized trial of the PhoenixCare intervention: program of case management and coordinated care for the seriously chronically ill. *J Palliat Med* 2006;9(1):111-126.
6. Jordhøy MS, Fayers P, Saltnes T, Ahlner-Elmqvist M, Jannert M, Kaasa S. A palliative-care intervention and death at home: a cluster randomized trial. *Lancet* 2000;356(9233):888-893.
7. Raftery JP, Addington-Hall JM, MacDonald LD, Anderson HR, Bland JM, Chamberlain J, Freeling P. A randomized controlled trial of the cost-effectiveness of a district co-ordinating service for terminally ill cancer patients. *Palliat Med* 1996;10(2):151-61.

Domain 1: Patient Care

Guideline 4 – Holistic Assessment and On-going Care Planning

Holistic assessment and on-going care planning are implemented in an interdisciplinary manner to meet the changing needs and wishes of patients, caregivers and families.

Rationale Patients approaching the end of life should receive an initial holistic assessment that covers their physical, social, spiritual and cultural needs and preferences. Identified needs and preferences can then be supported by members of the interdisciplinary team.¹⁻⁵ Subsequent on-going assessment and care planning should be proactive and responsive to patients’ changing needs.

Indicators

All	4.1	All patients identified as approaching the end of life undergo documented holistic assessments that cover the patient and family's physical, psychological, social, spiritual and cultural needs and preferences.
	4.2	There is evidence of individualised care plans made after the holistic assessment of the needs of patients, caregivers and families.
B + C	4.3	Assessment and care-planning reflect an interdisciplinary approach.
	4.4	There is evidence of on-going assessment and care planning at appropriate intervals that documents changes in the patient and family's needs, and response to treatment over time.
	4.5	Regular assessment of physical symptoms, and psychological and spiritual needs is conducted with the use of assessment tools where appropriate.
	4.6	There are mechanisms in place to identify and assess risks of self-harm.

Tools

1	Patient assessment forms: a. Forms and other resources from US providers, assembled by the Center to Advance Palliative Care ⁶ b. Forms from local Class C providers	- Annex A
2	Symptom assessment tools: a. Edmonton Symptom Assessment System (ESAS) ⁷⁻⁹ b. Palliative Care Outcomes Collaboration (PCOC) ¹⁰⁻¹²	-
3	Spiritual assessment tool: a. FICA Spiritual Assessment Tool ¹³⁻¹⁵ b. “Are you at peace?” One item to probe spiritual concerns at the End of Life ¹⁶ c. HOPE Spiritual Assessment Tool ¹⁷	-

References:

- Higginson IJ, Evans CJ. What is the evidence that palliative care teams improve outcomes for cancer patients and their families? *Cancer J* 2010;16(5):423-435.

2. Bakitas M, Lyons KD, Hegel MT, Balan S, Brokaw FC, Seville J, Hull JG, Li Z, Tosteson TD, Byock IR, Ahles TA. Effects of a palliative care intervention on clinical outcomes in patients with advanced cancer: the Project ENABLE II randomized controlled trial. *JAMA* 2009;302(7):741-9.3.
3. Puchalski C, Ferrell B, Virani R, Otis-Green S, Baird P, Bull J, Chochinov H, Handzo G, Nelson-Becker H, Prince-Paul M, Pugliese K, Sulmasy D. Improving the quality of spiritual care as a dimension of palliative care: The report of the consensus conference. *J Pall Med* 2009;12(10):885-904.
4. Abernethy AP, Currow DC, Fazekas BS, Luszcz MA, Wheeler JL, Kuchibhatla M. Specialized palliative care services are associated with improved short- and long-term caregiver outcomes. *Support Care Cancer* 2008;16(6):585-97.
5. Brumley R, Enguidanos S, Jamison P, Seitz R, Morgenstern N, Saito S, McIlwane J, Hillary K, Gonzalez J. Increased satisfaction with care and lower costs: results of a randomized trial of in-home palliative care. *J Am Geriatr Soc* 2007;55(7):993-1000.
6. Center to Advance Palliative Care (CAPC) Clinical Tools for Palliative Care Programs. Available online. <http://www.capc.org/tools-for-palliative-care-programs/clinical-tools/>. Last accessed 30th December 2013.
7. Edmonton Symptom Assessment System Revised. Available online <http://www.palliative.org/newpc/professionals/tools/esas.html>. Last accessed 30th December 2013.
8. Watanabe SM, Nikolaichuk CL, Beaumont C. The Edmonton Symptom Assessment System, a proposed tool for distress screening in cancer patients: development and refinement. *Psychooncology* 2012;21(9):977-85.
9. Watanabe SM, Nikolaichuk C, Beaumont C, Johnson L, Myers J, Strasser F. A multicenter study comparing two numerical versions of the Edmonton Symptom Assessment System in palliative care patients. *J Pain Symptom Manage* 2011;41(2):456-68.
10. Palliative Care Outcomes Collaboration (PCOC) Version 3 Dataset Forms. Available online <http://www.pcoc.org.au/>. Last accessed 30th December 2013.
11. Eagar K, Watters P, Currow DC, Aoun SM, Yates P. The Australian Palliative Care Outcomes Collaboration (PCOC)--measuring the quality and outcomes of palliative care on a routine basis. *Aust Health Rev* 2010;34(2):186-92.
12. Currow DC, Eagar K, Aoun S, Fildes D, Yates P, Kristjanson LJ. Is it feasible and desirable to collect voluntarily quality and outcome data nationally in palliative oncology care? *J Clin Oncol* 2008;26(23):3853-9.
13. FICA Spiritual Assessment Tool. Available online <http://smhs.gwu.edu/gwish/clinical/fica>. Last accessed 30th December 2013.
14. Puchalski CM. Integrating spirituality into patient care: an essential element of person-centered care. *Pol Arch Med Wewn* 2013;123(9):491-7.
15. Borneman T, Ferrell B, Puchalski CM. Evaluation of the FICA Tool for Spiritual Assessment. *J Pain Symptom Manage* 2010;40(2):163-73.
16. Steinhäuser KE, Voils CI, Clipp EC, Bosworth HB, Christakis NA, Tulsky JA. "Are You at Peace?" One Item to Probe Spiritual Concerns at the End of Life. *Arch Intern Med* 2006;166:101-105
17. Anandarajah G, Hight E. Spirituality and Medical Practice: Using the HOPE Questions as a Practical Tool for Spiritual Assessment. *Am Fam Physician* 2001;63(1):81-89

Domain 1: Patient Care

Guideline 5 – Advance Care Planning

All patients at the end of life have access to Advance Care Planning (ACP).

Rationale ACP helps to ensure that patients’ wishes are respected in the event that they become incapable of participating in treatment decisions, and allows for treatment at the end-of-life to be consistent with the patients’ preferences.¹⁻⁷

Indicators

All	5.1	There are systems in place to provide patients with life-limiting illness with information about, and access to, Advance Care Planning.
B + C	5.2	The service provider routinely conducts Advance Care Planning to ascertain and document patients’ and families’ preferences about treatment at the end-of-life, and fulfils these preferences as far as possible.
	5.3	There are systems in place to monitor if patients’ Advance Care Plans are honoured.

Tools

1	Informative, educational and publicity materials for healthcare staff and patients: a. Living Matters resource site ⁸ (including brochures, FAQs, ACP form templates and a Do-It-Yourself ACP workbook)	-
2	Examples of current referral procedures to trained ACP Facilitators within restructured hospitals	Annex B

References:

1. R Stein RA, Sharpe L, Bell ML, Boyle FM, Dunn SM, Clarke SJ. Randomized controlled trial of a structured intervention to facilitate end-of-life decision making in patients with advanced cancer. *J Clin Oncol* 2013;31(27):3403-10.
2. Epstein AS, Volandes AE, O'Reilly EM. Building on Individual, State, and Federal Initiatives for Advance Care Planning, an Integral Component of Palliative and End-of-Life Cancer Care. *J Oncol Pract* 2011;7(6):355-9.
3. Detering KM, Hancock AD, Reade MC, Silvester W. The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ* 2010;340:c1345.
4. Hammes BJ, Rooney BL, Gundram JD. A comparative, retrospective, observational study of the prevalence, availability, and specificity of advance care plans in a county that implemented an advance care planning microsystem. *J Am Geriatr Soc* 2010;58(7):1249-1255.
5. Teno JM, Gruneir A, Schwartz Z, Nanda A, Wetle T. Association between advance directives and quality of end-of-life care: a national study. *J Am Geriatr Soc* 2007;55(2):189-194.
6. Silveira MJ, Kim SY, Langa KM. Advance Directives and outcomes of surrogate Decision Making before Death. *N Engl J Med* 2010;362(13):1211-1218.
7. Zhang B, Wright AA, Huskamp HA, Nilsson ME, Maciejewski ML, Earle CC, Block SD. Health care costs in the last week of life: associations with end-of-life conversations. *Arch Intern Med* 2009;169:480-8.
8. Living Matters. Available online. <http://livingmatters.sg>. Last accessed 27th May 2014.

Domain 1: Patient Care

Guideline 6 – Patient-Centred Care

Patients receive care that is customized to their unique needs and preferences as informed by holistic assessments.

Rationale

Patients have unique needs and preferences, which may differ depending on their cultural background. The patient's quality of life is improved by care that is customised to their unique physical, emotional, cultural and spiritual needs.¹⁻⁴

Indicators

All	6.1	Each patient's unique cultural and spiritual needs are taken into consideration in the provision of care services.
	6.2	Information on the patient's condition is openly and sensitively communicated to the patient and family on a regular basis.
C	6.3	Spiritual support and therapy services (e.g. occupational, physical and speech therapy) are made available to patients when needed. Where these competencies do not exist within the service provider, there should be defined links to access these services.
	6.4	In the presence of complex ethical dilemmas beyond the resourced ability of service provider, there are policies and procedures to ensure access to ethics committees (or equivalent).

Tools

1	<p>Patient assessment forms:</p> <ul style="list-style-type: none"> a. Forms and other resources from US providers, assembled by the Center to Advance Palliative Care⁵ b. Forms from local Class C providers 	- Annex A
2	<p>Reference material for healthcare staff:</p> <ul style="list-style-type: none"> a. Online casebook on "Making Difficult Decisions with Patients and Families", developed by the NUS Centre for Biomedical Ethics⁶ 	

References:

1. Ellis PM. The importance of multidisciplinary team management of patients with non-small-cell lung cancer. *Curr Oncol* 2012;19(Suppl 1):S7-S15.
2. Janssen DJ, Spruit MA, Alsemgeest TP, Does JD, Schols JM, Wouters EF. A patient-centred interdisciplinary palliative care programme for end-stage chronic respiratory diseases. *Int J Palliat Nurs* 2010;16(4):189-94.
3. Mitchell GK, Del Mar CB, O'Rourke PK, Clavarino AM. Do case conferences between general practitioners and specialist palliative care services improve quality of life? A randomised controlled trial (ISRCTN 52269003). *Palliat Med* 2008;22(8):904-12.
4. Lloyd-Williams M, Reeve J, Kissane D. Distress in palliative care patients: developing patient-centred approaches to clinical management. *Eur J Cancer* 2008;44(8):1133-8.
5. Center to Advance Palliative Care (CAPC) Clinical Tools for Palliative Care Programs. Available online. <http://www.capc.org/tools-for-palliative-care-programs/clinical-tools/>. Last accessed 30th December 2013.

6. NUS Centre for Biomedical Ethics. Making Difficult Decisions with Patients and Families: A Singapore Casebook. Available online. <http://www.bioethicscasebook.sg/>. Last accessed 6th April 2014.

Domain 1: Patient Care

Guideline 7 – Care in the Last Days of Life

Care is taken to fulfil the needs of patients in the last days of life, as well as that of their caregivers and families.

Rationale

During the last hours and days of life the unique needs of patients and families should be taken into consideration, the comfort of patients maximised and their dignity respected.¹⁻⁸

Indicators

All	7.1	There is recognition and documentation of the patient's transition to the active dying phase, and communication to the patient, family and staff on the patient's imminent death.
	7.2	The family is educated on a timely basis on the signs and symptoms of imminent death in an age-appropriate, developmentally appropriate and culturally appropriate manner.
	7.3	Symptoms at the end of life are assessed and controlled, with referral to palliative care services if necessary.
	7.4	There is evidence of a plan in place to maximise patient comfort during the active dying phase and to support the family and caregivers.

Tools

1	<p>Guidebooks for staff:</p> <p>a. The Bedside Palliative Medicine Handbook,⁹ a practical guide to palliative medicine in Singapore developed by Tan Tock Seng Hospital</p>	-
2	<p>Education materials for patients and caregivers:</p> <p>a. Singapore Hospice Council's guide to terminal care¹⁰</p> <p>b. Patient/ caregiver information guides/booklets from local Class C providers</p>	Annex C

References:

1. Kehl KA, Kowalkowski JA. A systematic review of the prevalence of signs of impending death and symptoms in the last 2 weeks of life. *Am J Hosp Palliat Care* 2013;30(6):601-16.
2. Lundquist G, Rasmussen BH, Axelsson B. Information of Imminent Death or Not: Does It Make a Difference? *J Clin Onc* 2011;29:3927-3931.
3. Sy Heath JA, Clarke NE, Donath SM, McCarthy M, Anderson VA, Wolfe J. Symptoms and suffering at the end of life in children with cancer: an Australian perspective. *Med J Aust* 2010;192(2):71-5.
4. Murtagh FE, Addington-Hall J, Edmonds P, Donohoe P, Carey I, Jenkins K, Higginson IJ. Symptoms in the month before death for stage 5 chronic kidney disease patients managed without dialysis. *J Pain Symptom Manage* 2010;40(3):342-52.
5. Shinjo T, Morita T, Hirai K, Miyashita M, Sato K, Tsuneto S, Shima Y. Care for Imminently Dying Cancer Patients: Family Members' Experiences and Recommendations. *J Clin Onc* 2010;28:142-148.
6. Heath JA, Clarke NE, Donath SM, McCarthy M, Anderson VA, Wolfe J. Symptoms and suffering at the end of life in children with cancer: an Australian perspective. *Med J Aust* 2010;192(2):71-5.

7. Low JA, Pang WS, Lee A, Shaw RJ. A descriptive study of the demography, symptomology, management and outcome of the first 300 patients admitted to an independent hospice in Singapore. *Ann Acad Med Singapore* 1998;27(6):824-9
8. Tay WK, Shaw RJ, Goh CR. A survey of symptoms in hospice patients in Singapore. *Ann Acad Med Singapore* 1994;23(2):191-6.
9. Hum Allyn, Koh Mervyn (eds.). *The Bedside Palliative Medicine Handbook*. Singapore: Armour Publishing, 2013.
10. Singapore Hospice Council. A Guide to Terminal Care. Available online. http://www.singaporehospice.org.sg/9.2_guidetoterminalcare.htm. Last accessed 23rd January 2014.

Domain 2: Family and Caregiver Support

Guideline 8 – Caregiver Support

Caregivers of patients with life-limiting illness face significant stress in their roles, and their own practical and emotional needs need to be supported.

Rationale Caregivers of patients with life-limiting illness face significant stress in their roles. Studies have shown that caregiving may negatively impact on caregivers' health and work. There are also practical and emotional needs which need to be supported¹⁻¹¹ in order for them to be able to provide care more effectively.

Indicators

All	8.1	The patient's primary caregiver is identified at the initial assessment, and his or her needs are assessed and addressed on an on-going basis.
B + C	8.2	The primary caregiver is provided with education and training on their role, including strategies for self-care and coping with the demands of caregiving.

Tools

1	<p>Informative materials on resources for patients and caregivers:</p> <ul style="list-style-type: none"> a. Training <ul style="list-style-type: none"> i. Schedule of training sessions conducted by the HCA Palliative Caregivers Programme¹² b. Online resource and support portals for patients and caregivers <ul style="list-style-type: none"> i. awwa's Caregiver Handbook¹³ ii. Singapore Silver Pages, an online directory with eldercare information run by the Agency for Integrated Care¹⁴ iii. Singapore Hospice Council Online Resource for Patient and Caregivers¹⁵ iv. Online guide developed by Macmillan Cancer Support¹⁶ v. Online guide developed by National Hospice and Palliative Care Organisation (NHPCO)¹⁷ 	-
---	--	---

References:

1. Candy B, Jones L, Drake R, Leurent B, King M. Interventions for supporting informal caregivers of patients in the terminal phase of a disease. *Cochrane Database Syst Rev* 2011;6:CD007617.
2. Hudson PL, Remedios C, Thomas K. A systematic review of psychosocial interventions for family carers of palliative care patients. *BMC Palliat Care* 2010;9:17.
3. Northouse LL, Katapodi MC, Song L, Zhang L, Mood DW. Interventions with family caregivers of cancer patients: Meta-analysis of randomized trials. *CA: A Cancer Journal for clinicians* 2010;60:317-339.
4. Grande G, Stajduhar K, Aoun S, Toye C, Funk L, Addington-Hall J, Payne S, Todd C. Supporting lay carers in the end of life care: current gaps and future priorities. *Palliat Med* 2009;23(4):339-344.
5. Hudson P, Thomas T, Quinn K, Cockayne M, Braithwaite M. Teaching family carers about home-based palliative care: final results from a group education program. *J Pain Symptom Manage* 2009;38(2):299-308.
6. Docherty A, Owens A, Asadi-Lari M, Petchey R, Williams J, Carter YH. Knowledge and information needs of informal caregivers in palliative care: a qualitative systematic review. *Palliat Med* 2008;22(2):153-171.

7. Kwak J, Salmon JR, Acquaviva KD, Brandt K, Egan KA. Benefits of training family caregivers on experiences of closure during end-of-life care. *J Pain Symptom Manage* 2007;33(4):434-445.
8. Walsh K, Jones L, Tookman A, Mason C, McLoughlin J, Blizard R, King M. Reducing emotional distress in people caring for patients receiving specialist palliative care. Randomized trial. *Br J Psychiatry* 2007;190:142-147.
9. McMillan SC, Small BJ, Weitzner M, Schonwetter R, Tittle M, Moody L, Haley WE. Impact of coping skills intervention with family caregivers of hospice patients with cancer: a randomized clinical trial. *Cancer* 2006;106(1):214-222.
10. Harding R, Higginson IJ. What is the best way to help caregivers in cancer and palliative care? A systematic literature review of interventions and their effectiveness. *Palliat Med* 2003;17(1):63-74.
11. The Survey on Informal Caregiving summary report for MCYS. Available online. <http://app.msf.gov.sg/Publications/TheSurveyonInformalCaregiving.aspx>. Last accessed 31st March 2014.
12. Palliative Caregivers Programme. Available online. <http://hca.org.sg/services/palliative-caregivers-programme>. Last accessed 23rd January 2014.
13. awwa Caregiver's Handbook. Available online. http://www.awwa.org.sg/index.php?option=com_content&view=article&id=13&Itemid=23. Last accessed 12th February 2014.
14. Singapore Silver Pages. Available online. <https://www.silverpages.sg/>. Last accessed 12th February 2014.
15. Singapore Hospice Council Homepage. For patients and caregivers. Available online. <http://www.singaporehospice.org.sg/index.htm>. Last accessed 23rd January 2014.
16. Macmillan Cancer Support. Available online. <http://www.macmillan.org.uk/Home.aspx>. Last accessed 23rd January 2014.
17. Caring Connections. Available online. <http://www.caringinfo.org/>. Last accessed 23rd January 2014.

Domain 2: Family and Caregiver Support

Guideline 9 – Bereavement Care

Family members affected by a death are offered timely bereavement support appropriate to their needs and preferences.

Rationale

There should be timely identification of complications in grief experienced by families before and after the patient's death. The provision of direct bereavement support, or referral of families to bereavement services should be based on the assessed needs of the families¹⁻¹¹. Psychotherapeutic interventions have been found benefit those who have marked difficulties adjusting to the loss¹².

Indicators

All	9.1	Appropriate information about practical death-related issues (e.g. funeral arrangements) should be available when requested.
	9.2	Families and caregivers identified to be at risk of complicated grief are referred to bereavement support services.
B + C	9.3	There is evidence of a system in place to screen caregivers and families for bereavement needs, and to provide direct bereavement support or referral to bereavement support services where necessary.

Tools

1	Informative materials on bereavement support services:	
	<ul style="list-style-type: none"> a. Non-exhaustive list of bereavement support services in Singapore b. Pamphlets on grief and bereavement for families and caregivers from local Class C providers 	Annex D Annex E
2	Informative materials on funeral services/arrangements upon death:	
	<ul style="list-style-type: none"> a. NEA online guide¹³ on death registration, funeral arrangements and other practical death-related issues 	-
3	Checklist for staff for after-death procedures:	
	<ul style="list-style-type: none"> a. Examples from local Class C providers 	Annex F
4	Inventory of Complicated Grief-Revised ¹⁴	-

References:

1. Allen JY, Haley WE, Small BJ, Schonwetter RS, McMillan SC. Bereavement among hospice caregivers of cancer patients one year following loss: predictors of grief, complicated grief, and symptoms of depression. *J Palliat Med* 2013;16(7):745-51.
2. Hudson P, Remedios C, Zordan R, Thomas K, Clifton D, Crewdson M, Hall C, Trauer T, Bolleter A, Clarke DM, Bauld C. Guidelines for the psychosocial and bereavement support of family caregivers of palliative care patients. *J Palliat Med* 2012;15(6):696-702.
3. Widera EW, Block SD. Managing grief and depression at the end of life. *Am Fam Physician* 2012;86(3):259-64.
4. Williams AL, McCorkle R. Cancer family caregivers during the palliative, hospice, and bereavement phases: a review of the descriptive psychosocial literature. *Palliat Support Care* 2011;9(3):315-25.

5. Holtslander LF. Caring for bereaved family caregivers: analyzing the context of care. *Clin J Oncol Nurs*. 2008;12(3):501-6
6. Grassi L. Bereavement in families with relatives dying of cancer. *Curr Opin Support Palliat Care* 2007;1(1):43-9.
7. Stroebe M, Schut H, Stroebe W. Health outcomes of bereavement. *Lancet* 2007;370(9603):1960-73.
8. Kreicbergs UC, Lannen P, Onelov E, Wolfe J. Parental grief after losing a child to cancer: impact of professional and social support on long-term outcomes. *J Clin Oncol* 2007;25(22):3307-12.
9. Zhang B, El-Jawahri A, Prigerson HG. Update on bereavement research: evidence-based guidelines for the diagnosis and treatment of complicated bereavement. *J Palliat Med* 2006;9(5):1188-203.
10. Hudson PL. How well do family caregivers cope after caring for a relative with advanced disease and how can health professionals enhance their support? *J Palliat Med* 2006;9(3):694-703.
11. Shear K, Frank E, Houck PR, Reynolds CF 3rd. Treatment of complicated grief: a randomized controlled trial. *JAMA* 2005;293(21):2601-8.
12. Currier JM, Neimeyer RA, Berman JS. The Effectiveness of Psychotherapeutic Interventions for Bereaved Persons: A Comprehensive Quantitative Review. *Psychological Bulletin* 2008; 134(5):648–661.
13. National Environment Agency. When a Loved One Passes Away. Available online. <http://www.nea.gov.sg/passesaway/>. Last accessed 12th February 2014.
14. Prigerson HG, Maciejewski PK, Reynolds CF, III, Bierhals AJ, Newsom JT, Fasiczka A, Frank E, Doman J, Miller M. The inventory of complicated grief: a scale to measure maladaptive symptoms of loss. *Psychiatry Research* 1995;59(1-2): 65-79.

Domain 3: Staff and Volunteer Management

Guideline 10 – Qualified Staff and Volunteers

Care for those approaching the end of life is delivered by staff and volunteers (where applicable) with the appropriate qualifications and skill mix for the level of service offered, and who demonstrate on-going participation in training and development.

Rationale

A competent and adequate workforce is key to the provision of quality care. There should be an adequate team of healthcare professionals with the appropriate training to meet the needs of patients at the end of life.¹⁻⁵ As service providers may have different service models, they should customise staffing norms according to the roles and responsibilities of the staff. Volunteers may complement the healthcare team in the provision of care, and must receive the appropriate training to perform their duties effectively.

Indicators

All	10.1	All staff and volunteers demonstrate evidence of palliative care training appropriate to the level of care provided.
C	10.2	Palliative care teams consist of an interdisciplinary team of skilled palliative care professionals, including physicians, nurses and social workers and/or counsellors/pastoral staff. Where these competencies do not exist within a team, there should be clearly defined links to access these through a service level agreement or similar.
	10.3	Staffing levels are adequate to deliver the care needed by patients.
	10.4	Palliative care professionals are appropriately trained, credentialed and/or certified in their area of expertise.
	10.5	Where volunteers are part of the team, there is evidence of policies in place to ensure proper screening, recruitment and on-going training of volunteers.

Tools

1	Informative materials for staff on relevant training: a. List of courses available at the AIC Learning Institute ⁶ b. List of training available in Singapore	- Annex I
2	Recommended staffing norms: a. Inpatient hospice: 2011 National Strategy for Palliative Care (NSPC) ⁷ b. Home and day hospice care: 2009 Singapore Hospice Council Home Care and Day Hospice Services Workgroup Report	- Annex J

References:

1. Henoch I, Danielson E, Strang S, Browall M, Melin-Johansson C. Training Intervention for Health Care Staff in the Provision of Existential Support to Patients With Cancer: A Randomized, Controlled Study.
2. J Pain Symptom Manage 2013;46(6):785-94Kang J, Kim Y, Yoo YS, Choi JY, Koh SJ, Jho HJ, Choi YS, Park J, Moon do H, Kim do Y, Jung Y, Kim WC, Lim SH, Hwang SJ, Choe SO, Jones D. Developing competencies for multidisciplinary hospice and palliative care professionals in Korea. Support Care Cancer. 2013;21(10):2707-17.

3. McCabe MR, Goldhammer D, Mellor D, Hallford D, Davison T. Evaluation of a training program to assist care staff to better recognize and manage depression among palliative care patients and their families. *J Palliat Care* 2012;28(2):75-82.
4. Juba KM. Pharmacist credentialing in pain management and palliative care. *J Pharm Pract*. 2012;25(5):517-20
5. Wittenberg-lyles E, Schneider G, Oliver DP. Results from the national hospice volunteer training survey. *J Pall Med* 2010;13(3):261-265.
6. AIC Learning Institute. Available online. <http://www.aic.sg/learninginstitute/>. Last accessed 23rd January 2014.
7. National Strategy for Palliative Care. Available online. http://www.duke-nus.edu.sg/sites/default/files/Report_on_National_Strategy_for_Palliative_Care%20Jan2012.pdf. Last accessed 23rd January 2014.

Domain 3: Staff and Volunteer Management

Guideline 11 – Staff and Volunteer Self-Care

Staff and volunteers reflect on practice, maintain effective self-care strategies and have access to support in dealing with the psychological stress associated with working among the terminally ill and bereaved.

Rationale The care of patients near the end of life and the support of their family members may have an emotional and spiritual toll on healthcare workers and volunteers.¹⁻⁶ The ability to reflect on their practice and opportunities to express their feelings related to interactions with patients and their families should be encouraged as part of the culture of service providers caring for patients near the end-of-life.⁷⁻¹⁰

Indicators

B + C	11.1	There are strategies in place to provide situational support, critical incident debriefing and response.
	11.2	Education is provided to help staff and volunteers develop effective coping strategies.
C	11.3	Staff have access to confidential employee assistance programs and/or counselling services.

Tools

-	-	-
---	---	---

References:

1. Chochinov HM. Dignity in care: time to take action. *J Pain Symptom Manage* 2013;46(5):756-9.
2. Mougalian SS, Lessen DS, Levine RL, Panagopoulos G, Von Roenn JH, Arnold RM, Block SD, Buss MK. Palliative care training and associations with burnout in oncology fellows. *J Support Oncol* 2013;11(2):95-102.
3. Phillips J, Andrews L, Hickman L. Role Ambiguity, Role Conflict, or Burnout: Are These Areas of Concern for Australian Palliative Care Volunteers? Pilot Study Results. *Am J Hosp Palliat Care* 2013 Oct 3. [Epub ahead of print]
4. Peters L, Cant R, Sellick K, O'Connor M, Lee S, Burney S, Karimi L. Is work stress in palliative care nurses a cause for concern? A literature review. *Int J Palliat Nurs* 2012;18(11):561-7.
5. Claxton-Oldfield S, Claxton-Oldfield J. Should I stay or should I go: a study of hospice palliative care volunteer satisfaction and retention. *Am J Hosp Palliat Care* 2012;29(7):525-30.
6. Pereira SM, Fonseca AM, Carvalho AS. Burnout in Palliative Care: A systematic review. *Nurs Ethics*. 2011;18(3):317-26.
7. Sanchez-Reilly S, Morrison LJ, Carey E, Bernacki R, O'Neill L, Kapo J, Periyakoil VS, Thomas Jde L. Caring for oneself to care for others: physicians and their self-care. *J Support Oncol* 2013;11(2):75-81.
8. Melo CG, Oliver D. Can addressing death anxiety reduce health care workers' burnout and improve patient care? *J Palliat Care* 2011;27(4):287-95.
9. Swetz KM, Harrington SE, Matsuyama RK, Shanafelt TD, Lyckholm LJ. Strategies for avoiding burnout in hospice and palliative medicine: peer advice for physicians on achieving longevity and fulfillment. *J Palliat Med* 2009;12(9):773-7.
10. Kearney MK, Weininger RB, Vachon ML, Harrison RL, Mount BM. Self-care of physicians caring for patients at the end of life: "Being connected... a key to my survival". *JAMA* 2009;301(11):1155-64.

Domain 4: Safe Care

Guideline 12 – Access to and Use of Opioids

Patients at the end of life should have access to opioids for symptom control, with guidelines and processes in place to ensure safe and effective use.

Rationale
Opioids are commonly used medications in palliative care. While useful, there are also risks and unwanted effects associated with unregulated use.¹⁻⁵ Education and guidelines are necessary to ensure optimal symptom control with minimal side effects.⁶⁻¹¹

Indicators

All	12.1	Service providers caring for patients at the end of life have access to opioids.
	12.2	The use of opioids in the management of symptoms should be directed by evidence, driven by need, and administered under guidelines with appropriate monitoring.
	12.3	There is evidence of patient and caregiver education on the safe use of opioids.
	12.4	There is adherence to the legal requirements of the Misuse of Drugs Act ¹⁰ on the prescription, safe storage, dispensing, administration, disposal and report of errors in the use of opioids.

Tools

1	Reference charts/ guidebooks for clinical staff: <ul style="list-style-type: none"> a. Opioid conversion charts from local Class C providers b. The Bedside Palliative Medicine Handbook,¹⁰ a practical guide to palliative medicine in Singapore developed by Tan Tock Seng Hospital¹¹ c. NICE guidelines 2012. Opioids in Palliative Care: safe and effective prescribing of strong opioids for pain in palliative care for adults.¹² 	Annex G - -
2	Educational materials for patients: <ul style="list-style-type: none"> a. Singapore Hospice Council pamphlet on using morphine 	Annex H

References:

1. Dowell D, Kunins HV, Farley TA. Opioid analgesics — risky drugs, not risky patients. *JAMA* 2013;309(21):2219-2220.
2. Boyer EW. Management of opioid analgesic overdose. *N Engl J Med* 2012;367(2):146-55.
3. Schisler RE, Groninger H, Rosielle DA. Counseling patients on side effects and driving when starting opioids #248. *J Palliat Med* 2012;15(4):484-5.
4. Webster PC. Medically induced opioid addiction reaching alarming levels. *CMAJ* 2012;184(3):285-6.
5. Kurita GP, Sjøgren P, Ekholm O, Kaasa S, Loge JH, Poviloniene I, Klepstad P. Prevalence and predictors of cognitive dysfunction in opioid-treated patients with cancer: A multinational study. *J Clin Oncol* 2011;29(10):1297-3032.
6. Ho KY, Chua NH, George JM, Yeo SN, Main NB, Choo CY, Tan JW, Tan KH, Ng BY; Pain Association of Singapore Task Force. Evidence-based guidelines on the use of opioids in chronic non-cancer pain--a

consensus statement by the Pain Association of Singapore Task Force. *Ann Acad Med Singapore*. 2013;42(3):138-52

7. Caraceni A, Hanks G, Kaasa S, Bennett MI, Brunelli C, Cherny N, Dale O, De Conno F, Fallon M, Hanna M, Haugen DF, Juhl G, King S, Klepstad P, Laugsand EA, Maltoni M, Mercadante S, Nabal M, Pigni A, Radbruch L, Reid C, Sjogren P, Stone PC, Tassinari D, Zeppetella G; European Palliative Care Research Collaborative (EPCRC); European Association for Palliative Care (EAPC). Use of opioid analgesics in the treatment of cancer pain: evidence-based recommendations from the EAPC. *Lancet Oncol* 2012;13(2):e58-68
8. Sheinfeld Gorin S, Krebs P, Badr H, Janke EA, Jim HS, Spring B, Mohr DC, Berendsen MA, Jacobsen PB. Meta-analysis of psychosocial interventions to reduce pain in patients with cancer. *J Clin Oncol* 2012;30(5):539-47.
9. Bennett MI, Bagnall AM, José Closs S. How effective are patient-based educational interventions in the management of cancer pain? Systematic review and meta-analysis. *Pain* 2009;143(3):192-9.
10. Attorney-General's Chambers. Misuse of Drugs Act. Available online <http://statutes.agc.gov.sg/aol/search/display/view.w3p?page=0;query=DocId%3A%22c13adadb-7d1b-45f8-a3bb-92175f83f4f5%22%20Status%3Apublished%20Depth%3A0;rec=0;whole=yes>. Last accessed 18 February 2014.
11. Hum Allyn, Koh Mervyn (eds.). *The Bedside Palliative Medicine Handbook*. Singapore: Armour Publishing, 2013.
12. NICE clinical guidelines 2012. Opioids in palliative care: safe and effective prescribing of strong opioids for pain in palliative care of adults. Available online. <http://www.nice.org.uk/nicemedia/live/13745/59285/59285.pdf>. Last accessed 23rd January 2014.

Domain 4: Safe Care

Guideline 13 – Clinical Quality Improvement

The service is committed to improvement in clinical and management practices.

Rationale

Palliative care services are committed to a high quality of care and support for all patients and their families. Services should strive to benchmark quality through regular and systematic measurement, analysis, review, evaluation, goal setting and revision of care processes.¹⁻¹⁸

Indicators

B + C	13.1	Presence of a framework to guide a program of quality improvement, that consists of: (a) Evidence of dissemination and incorporation of quality improvement (QI) findings into practice (b) Clinical and performance criteria consistent with professional standards (c) An on-going, proactive program for identifying and reducing unanticipated adverse events and safety risks to patients
	C	13.2 There is evidence of the implementation of quality improvement projects to support on-going service evaluation and development.
	13.3	There is evidence of on-going evaluation of patients' and families' satisfaction with care, and necessary measures to improve these.

Tools

1	Quality improvement and assurance framework: a. Examples from local Class C providers	Annex K
2	Resources for standardised and validated clinical assessment tools: a. Palliative Care Outcomes Collaboration (PCOC) Assessment Toolkit ¹⁹ b. Toolkit of Instruments to Measure End-of-Life Care (TIME) ²⁰	-

References:

- De Roo ML, Leemans K, Claessen SJ, Cohen J, Pasman HR, Deliens L, Francke AL; EURO IMPACT. Quality indicators for palliative care: update of a systematic review. *J Pain Symptom Manage* 2013;46(4):556-72.
- Woitha K, Van Beek K, Ahmed N, Jaspers B, Mollard JM, Ahmedzai SH, Hasselaar J, Menten J, Vissers K, Engels Y. Validation of quality indicators for the organization of palliative care: A modified RAND Delphi study in seven European countries (the Europall project). *Palliat Med* 2013 Jul 16. [Epub ahead of print]
- Leemans K, Cohen J, Francke AL, Vander Stichele R, Claessen SJ, Van den Block L, Deliens L. Towards a standardized method of developing quality indicators for palliative care: protocol of the Quality indicators for Palliative Care (Q-PAC) study. *BMC Palliat Care* 2013;12:6.
- Higashi T, Nakamura F, Saruki N, Sobue T. Establishing a quality measurement system for cancer care in Japan. *Jpn J Clin Oncol* 2013;43(3):225-32.

5. De Lima L, Bennett MI, Murray SA, Hudson P, Doyle D, Bruera E, Granda-Cameron C, Strasser F, Downing J, Wenk R. International Association for Hospice and Palliative Care (IAHPC) List of Essential Practices in Palliative Care. *J Pain Palliat Care Pharmacother* 2012;26(2):118-22.
6. Raijmakers N, Galushko M, Domeisen F, Beccaro M, Lundh Hagelin C, Lindqvist O, Popa-Velea O, Romotzky V, Schuler S, Ellershaw J, Ostgathe C; OPCARE9. Quality indicators for care of cancer patients in their last days of life: literature update and experts' evaluation. *J Palliat Med* 2012;15(3):308-16.
7. Agar M, Lockett T. Outcome measures for palliative care research. *Curr Opin Support Palliat Care* 2012;6(4):500-7.
8. Hanson LC, Rowe C, Wessell K, Caprio A, Winzelberg G, Beyea A, Bernard SA. Measuring palliative care quality for seriously ill hospitalized patients. *J Pall Med* 2012;15(7):798-804.
9. Penrod JD, Pronovost PJ, Livote EE, et al. Meeting standards of high-quality intensive unit palliative care: Clinical performance and predictors. *Critical Care Med* 2012;40(4):1105-1112.
10. El-Jawahri A, Greer JA, Temel JS. Does palliative care improve outcomes for patients with incurable illness? A review of the evidence. *J Support Oncol* 2011;9(3):87-94.
11. Neuss M, Gilmore TR, Kadlubek P. Tools for measuring and improving the quality of oncology care: the Quality Oncology Practice Initiative (QOPI) and the QOPI certification program. *Oncology (Williston Park)* 2011;25(10):880, 883, 886-7.
12. Weissman DE, Morrison RS, Meier DE. Center to Advance Palliative Care palliative care clinical care and customer satisfaction metrics consensus recommendations. *J Palliat Med* 2010;13(2):179-84.
13. Schenck AP, Rokoske FS, Durham DD, Cagle JG, Hanson LC. The PEACE project: Identification of quality measures for hospice and palliative care. *J Pall Med* 2010;13(12):1451-1459.
14. Walling AM, Asch SM, Lorenz KA, Roth CP, Barry T, Kahn KL, Wenger NS. The quality of care provided to hospitalized patients at the end of life. *Arch Int Med* 2010;170(12):1057-1063.
15. Weissman DE, Meier DE. Center to Advance Palliative Care inpatient unit operational metrics: consensus recommendations. *J Palliat Med* 2009;12(1):21-5.
16. Weissman DE, Meier DE, Spragens LH. Center to Advance Palliative Care palliative care consultation service metrics: consensus recommendations. *J Palliat Med* 2008;11(10):1294-8.
17. Ferrell B, Connor SR, Cordes A, Dahlin CM, Fine PG, Hutton N, Leenay M, Lentz J, Person JL, Meier DE, Zuroski K; National Consensus Project for Quality Palliative Care Task Force Members. The national agenda for quality palliative care: the National Consensus Project and the National Quality Forum. *J Pain Symptom Manage* 2007;33(6):737-44.
18. Twaddle ML, Maxwell TL, Cassel JB, Liao S, Coyne PJ, Usher BM, Amin A, Cuny J. Palliative care benchmarks from academic medical centers. *J Palliat Med* 2007;10:86-98
19. Palliative Care Outcomes Collaboration. Available online. <http://www.pcoc.org.au/>. Last accessed 16th January 2014.
20. Toolkit of Instruments to Measure End-of-Life Care. Available online. <http://as800.chcr.brown.edu/pcoc/>. Last accessed 23rd January 2014.

Acknowledgements

This set of Guidelines was developed by the Standards Development Subgroup of the National Strategy for Palliative Care Implementation Taskforce:

Dr Angel Lee (Chairman)	Senior Consultant, Dover Park Hospice Senior Consultant, Department of Palliative Medicine, Tan Tock Seng Hospital
Dr Patricia Neo (Vice-Chairman)	Consultant, Department of Palliative Medicine, National Cancer Centre Singapore
Dr R. Akhileswaran	CEO and Medical Director, HCA Hospice Care Chairman, Singapore Hospice Council
Ms Chen Wei Ting	Advanced Practice Nurse, Tan Tock Seng Hospital
Ms Ng Tzer Wee	Manager and Principal Medical Social Worker, Tan Tock Seng Hospital
Dr Ong Yew Jin	Consultant, Assisi Hospice
A/Prof Edward Poon	Director of Nursing, Ang Mo Kio – Thye Hua Kwan Hospital
Dr Angeline Seah	Senior Consultant, Department of Geriatric Medicine, Khoo Teck Puat Hospital
Dr Siew Chee Weng	Family Physician Director, ElderPrime Medical Pte Ltd

In consultation with:

A/Prof John Abisheganaden	Senior Consultant, Department of Respiratory Medicine, Tan Tock Seng Hospital
Dr Noreen Chan	Senior Consultant, Department of Haematology-Oncology, National University Hospital
Ms Chee Wai Yee	Head, Allied Health Services, Dover Park Hospice
Dr Irwin Chung	Director, Care Integration Division and Consultant, Regional Health System & Primary Care Development Division, Agency for Integrated Care
A/Prof Cynthia Goh	Deputy Chairperson, Lien Centre for Palliative Care Senior Consultant, Department of Palliative Medicine, National Cancer Centre Singapore
Dr Shirlynn Ho	Associate Consultant, Department of Palliative Medicine, National Cancer Centre Singapore
Dr Koh Lip Hoe	Consultant, Department of Geriatric Medicine, Changi General Hospital
Dr Kok Jaan-Yang	Senior Consultant, Palliative Care, Parkway Cancer Centre
Dr Lee Kheng Hock	Head of Department, Senior Consultant, Singapore General Hospital
Mrs Lee Lay Beng	Principal Medical Social Worker, Tan Tock Seng Hospital
Ms Low Mui Lang	Executive Director, The Salvation Army Peacehaven Nursing Home
Dr Rina Nga	Resident Physician, Hospice Services Department, Singapore Cancer Society
A/Prof Pang Weng Sun	Chairman, Medical Board, Yishun Community Hospital
Dr Pek Wee Yang	Head, Department of Medicine, Khoo Teck Puat Hospital
Dr David Sim	Consultant, National Heart Centre
Sr Geraldine Tan	Administrator, St Joseph's Home
Dr Tan Yew Seng	Medical Director, Assisi Hospice

Mr Ivan Woo Mun Hong	Principal Medical Social Worker, Department of Care and Counselling, Tan Tock Seng Hospital
Dr Wu Huei Yaw	Medical Director, Dover Park Hospice
Dr Grace Yang	Associate Consultant, Department of Palliative Medicine, National Cancer Centre


Supported by:

Dr Winston Chin	Assistant Director, Ageing Planning Office, MOH
Ms Lilian Lee	Manager, Ageing Planning Office, MOH
Ms Lin Yongqing	Assistant Manager, Ageing Planning Office, MOH
Ms Katherine Soh	Principal Manager, Regulatory Policy and Legislation Division, MOH (formerly of Standards and Quality Improvement Division)

And is endorsed by:



Singapore Hospice Council



**Chapter of Palliative Medicine,
College of Physicians, Singapore**



**Palliative Care Nurses Chapter,
Singapore Nurses Association**



Singapore Association of Social Workers

